

FETAL DIAGNOSIS SERVICE

Referral Criteria

- adrenal anomaly
- ambiguous genitalia
- ascites
- bladder exstrophy
- bladder outlet obstruction
- bowel dilatation, intestinal obstruction
- brain anomalies: absent CSP, agenesis/dysgenesis of the corpus callosum, Chiari II malformation, dilated ventricles, holoprosencephaly, posterior fossa abnormality, etc.
- cardiac malformation, dextrocardia, rhabdomyoma
- cleft lip and/or palate
- club foot
- cystic hygroma (>17 weeks) (less than 17 weeks gestation should be referred to Medical Genetics)
- diaphragmatic hernia
- ectopic kidney
- ectrodactyly
- encephalocele
- hydronephrosis (>10mm), hydroureter
- hydrops (> 17wks) (less than 17 weeks should be referred to Medical Genetics)
- limb reduction defects
- meconium peritonitis
- megacystis
- micro/retrognathia
- microcephaly (HC < 1st percentile)
- multicystic or echogenic kidneys
- multiple pterygia, fetal akinesia
- myelomeningocele
- oligohydramnios (detected prior to 26 weeks gestation)
- omphalocele
- pentology of Cantrell
- pericardial effusion (> 3 mm)
- pleural effusion
- polydactyly
- polyhydramnios, severe (AFI \geq 350 mm)
- renal agenesis
- sacrococcygeal teratoma
- scoliosis, kyphosis
- skeletal dysplasia
- spina bifida
- subcutaneous edema
- talipes equinovarus
- ventriculomegaly (10 mm or greater)
- vertebral segmentation anomaly